

TOP-BESS MODEL AND ITS PHENOMENOLOGY

M. Gintner^{1,2}, J. Juráň², I. Melo¹

¹ U. of Žilina, Žilina, Slovakia

² IEAP CTU Prague, Czech Republic

PLANCK 2012 Warsaw
May 30, 2012

OUTLINE

1 INTRODUCTION

2 TOP-BESS MODEL

3 PHENOMENOLOGY

OUTLINE

1 INTRODUCTION


2 TOP-BESS MODEL

3 PHENOMENOLOGY

CHALLENGE



CHALLENGE




CHALLENGE



HIGGS
DOUBLET
???

BSM ???

- weakly interacting
- strongly interacting
- extra-dims



EFFECTIVE DESCRIPTION OF STRONG ESB

$SU(2)_L \times U(1)_Y$ broken *dynamically*:

- *not* solvable perturbatively
- *chiral effective Lagrangian* for **Goldstone bosons**
nonlinear sigma model
- ... + **resonances**
scalar, vector, ...

LHC → the *lightest* BSM resonances

OUTLINE

1 INTRODUCTION

2 TOP-BESS MODEL

3 PHENOMENOLOGY

BESS MODEL

BREAKING ELECTROWEAK SYMMETRY STRONGLY

- R. Casalbuoni *et al*, PLB155,95(1985); NPB282,235(1987)
- \mathcal{HSM} + new vector resonances
- Hidden Local Symmetry [M. Bando *et al* (1984)]
- *global symmetry:*

$$SU(2)_L \times SU(2)_R \times U(1)_{B-L} \times SU(2)_{HLS} \xrightarrow{SSB} SU(2)_{L+R} \times U(1)_{B-L}$$

- *local symmetry:*

$$\begin{array}{ccc} SU(2)_L \times U(1)_Y \times SU(2)_{HLS} & \xrightarrow{SSB} & U(1)_{em} \\ g & g' & g'' \\ & & e \end{array}$$

BESS LAGRANGIAN: ESB

ESB sector (NGB's + gauge bosons):

$$\mathcal{L}_{ESB} = \underbrace{-v^2 \text{Tr}[(\bar{\omega}_\mu^\perp)^2] - \alpha v^2 \text{Tr}[(\bar{\omega}_\mu^\parallel)^2]}_{\sim \text{gauged NLSM}} + \underbrace{\mathcal{L}_{GB}(W, B, V)}_{\text{kin. terms} + \text{self-interactions}}$$

GB masses + mixing



$$W^\pm, Z \quad A \quad V^\pm, V^0$$

$$M_V = \frac{1}{2}\sqrt{\alpha} g'' v$$

BESS LAGRANGIAN: FERMIONS

fermion sector (SM fermions):

$$\mathcal{L}_f^{BESS} = \underbrace{\mathcal{L}_f^{SM}(W, B)}_{\text{mixing induced } Vff \text{ cplngs}} + \underbrace{\mathcal{L}_f^{BSM}(W, B, V)}_{\text{direct universal chiral } Vff \text{ cplngs}}$$

$\sim 1/g''$

$\sim b_L g'' , \sim b_R g''$

universality:

- ν_R absence, K_L - K_S mass difference $\Rightarrow b_R \rightarrow 0$
- $Z \rightarrow \bar{f}f$ at LEP \Rightarrow tight limits on b_L

BESS LAGRANGIAN: FERMIONS

fermion sector (SM fermions):

$$\mathcal{L}_f^{BESS} = \underbrace{\mathcal{L}_f^{SM}(W, B)}_{\text{mixing induced } Vff \text{ cplngs}} + \underbrace{\mathcal{L}_f^{BSM}(W, B, V)}_{\text{direct universal chiral } Vff \text{ cplngs}}$$

$\sim 1/g''$ $\sim b_L g'' , \sim b_R g''$

universality:

- ν_R absence, K_L - K_S mass difference $\Rightarrow b_R \rightarrow 0$
- $Z \rightarrow \bar{f}f$ at LEP \Rightarrow tight limits on b_L

OUTSTANDING TOP QUARK

$m_t \approx v/\sqrt{2}$ → special role in ESB?

new physics behind m_t



ESB related

Extended TC, ...



ESB unrelated

Topcolor Assisted TC, ...

TOP-BESS MODEL

PHYS. REV. D84, 035013 (2011)

*ESB sector \equiv BESS**fermion sector (SM fermions):*

$$\mathcal{L}_f^{tBESS} = \underbrace{\mathcal{L}_f^{SM}(W, B)}_{\text{mixing induced } Vff \text{ cplngs}} + \underbrace{\mathcal{L}_{(t,b)}^{BSM}(W, B, V)}_{\text{direct chiral cplngs}} + \underbrace{\mathcal{L}_{(t,b)}^{BSM'}(W, B)}_{\text{new } W/B \text{ cplngs}}$$

*mixing induced
 Vff cplngs*

$\sim 1/g''$

direct chiral cplngs

$Vtt \dots$	$\sim b_{L,R} \cdot g''$
$Vt_L b_L \dots$	$\sim b_L \cdot g''$
$Vb_L b_L \dots$	$\sim b_L \cdot g''$
$Vt_R b_R \dots$	$\sim p b_R \cdot g''$
$Vb_R b_R \dots$	$\sim p^2 b_R \cdot g''$

$0 \leq p \leq 1$

new W/B cplngs

$Xtt \dots$	$\sim \lambda_{L,R} \cdot g_X$
$Xt_L b_L \dots$	$\sim \lambda_L \cdot g_X$
$Xb_L b_L \dots$	$\sim \lambda_L \cdot g_X$
$Xt_R b_R \dots$	$\sim p \lambda_R \cdot g_X$
$Xb_R b_R \dots$	$\sim p^2 \lambda_R \cdot g_X$

$X=W,B, \quad g_X = g, g'$

TOP-BESS MODEL

PHYS. REV. D84, 035013 (2011)

*ESB sector \equiv BESS**fermion sector (SM fermions):*

$$\mathcal{L}_f^{tBESS} = \underbrace{\mathcal{L}_f^{SM}(W, B)} + \underbrace{\mathcal{L}_{(t,b)}^{BSM}(W, B, V)} + \underbrace{\mathcal{L}_{(t,b)}^{BSM'}(W, B)}$$

*mixing induced
Vff cplngs*

$$\sim 1/g''$$

direct chiral cplngs

$Vtt \dots$	$\sim b_{L,R} \cdot g''$
$Vt_L b_L \dots$	$\sim b_L \cdot g''$
$Vb_L b_L \dots$	$\sim b_L \cdot g''$
$Vt_R b_R \dots$	$\sim p b_R \cdot g''$
$Vb_R b_R \dots$	$\sim p^2 b_R \cdot g''$

new W/B cplngs

$Xtt \dots$	$\sim \lambda_{L,R} \cdot g_X$
$Xt_L b_L \dots$	$\sim \lambda_L \cdot g_X$
$Xb_L b_L \dots$	$\sim \lambda_L \cdot g_X$
$Xt_R b_R \dots$	$\sim p \lambda_R \cdot g_X$
$Xb_R b_R \dots$	$\sim p^2 \lambda_R \cdot g_X$

$$0 \leq p \leq 1$$

$$X=W,B, \quad g_X = g, g'$$

OUTLINE


1 INTRODUCTION

2 TOP-BESS MODEL

3 PHENOMENOLOGY

V DECAY WIDTHS

- $V^0 \rightarrow W^+W^- + t\bar{t} + b\bar{b} + \dots$
- $V^+ \rightarrow W^+Z + t\bar{b} + \dots$
- $\Gamma \sim 10 - 100 \text{ GeV}$




effect of λ 's is negligible

UNITARITY & PERTURBATIVITY

Unitarity:

- GB scatterings:
 $W_L^+ W_L^-$, $Z_L Z_L$,
 $W_L^\pm Z_L$, $W_L^\pm W_L^\pm$
- tree level
- Equivalence Theorem



Perturbativity:

$$\frac{g''}{2} \lesssim 4\pi \quad \Rightarrow \quad g'' \lesssim 30$$

LOW-ENERGY LIMITS

LOW-ENERGY LAGRANGIAN

$$\dots \alpha \rightarrow \infty \Rightarrow M_V \rightarrow \infty$$

free parameters:

$$\begin{aligned}x &= g/g'' \\ \Delta L &= b_L - 2\lambda_L \\ \Delta R &= b_R + 2\lambda_R \\ p\end{aligned}$$


EXPERIMENT: LEP + SLC + TEVATRON

$$\begin{aligned}\epsilon_1 &\stackrel{\text{exp}}{=} (+5.4 \pm 1.0) \times 10^{-3} \\ \epsilon_2 &\stackrel{\text{exp}}{=} (-8.9 \pm 1.2) \times 10^{-3} \\ \epsilon_3 &\stackrel{\text{exp}}{=} (+5.34 \pm 0.94) \times 10^{-3} \\ \Gamma(Z \rightarrow \bar{b}b) &\stackrel{\text{exp}}{=} (0.3773 \pm 0.0013) \text{ GeV} \\ B.R.(B \rightarrow X_s \gamma) &\stackrel{\text{exp}}{=} (3.55 \pm 0.26) \times 10^{-4}\end{aligned}$$

LOW-ENERGY LIMITS FOR FERMION PARAMS

 χ^2 -FIT:

- 5 observables:
 $\epsilon_1, \epsilon_2, \epsilon_3, \Gamma(Z \rightarrow b\bar{b}),$
 $\text{BR}(B \rightarrow X_s\gamma)$
- 2 fitted params:
 $\Delta L = b_L - 2\lambda_L$
 $\Delta R = b_R + 2\lambda_R$
- fixed params:
 $g'' = 20$
 $p = 0, 0.5, 1$
 $\Lambda = 1 \text{ TeV}$
- confidence levels:
1 sigma ... 39%
2 sigma ... 86%
3 sigma ... 99%




THE DEATH VALLEY

direct + indirect *cplgs* \Rightarrow DV

THE DEATH VALLEY


direct + indirect *cplgs* \Rightarrow DV



$$g'' = 10, \lambda_L = \lambda_R = 0$$


The Death Valley regions of the $V \rightarrow t\bar{t}/b\bar{b}/tb$ decays.

THE DEATH VALLEY

direct + indirect *cplgs* \Rightarrow DV*The Death Valley regions of the $V \rightarrow t\bar{t}/b\bar{b}/tb$ decays.*

HIDING THE PEAK

$$M_V = 1 \text{ TeV}, \ g'' = 20, \ p = 0, \ \lambda_R = 0$$




- no direct cplng
 $b_L = 0$
 $b_R = 0$
 $\lambda_L = 0$
- outside the DV
 $b_L = -0.010$
 $b_R = +0.030$
 $\lambda_L = 0$
- $t\bar{b}$ & $b\bar{b}$ in the DV
 $b_L = +0.009$
 $b_R = +0.030$
 $\lambda_L = +0.006$
- all in the DV
 $b_L = +0.0098$
 $b_R = +0.0034$
 $\lambda_L = +0.006$



HIDING THE PEAK

$$M_V = 1 \text{ TeV}, \ g'' = 20, \ p = 0, \ \lambda_R = 0$$




- no direct cplng
 $b_L = 0$
 $b_R = 0$
 $\lambda_L = 0$
- outside the DV
 $b_L = -0.010$
 $b_R = +0.030$
 $\lambda_L = 0$
- $t\bar{b}$ & $b\bar{b}$ in the DV
 $b_L = +0.009$
 $b_R = +0.030$
 $\lambda_L = +0.006$
- all in the DV
 $b_L = +0.0098$
 $b_R = +0.0034$
 $\lambda_L = +0.006$



HIDING THE PEAK

$$M_V = 1 \text{ TeV}, \ g'' = 20, \ p = 0, \ \lambda_R = 0$$




- **no direct cplng**
 $b_L = 0$
 $b_R = 0$
 $\lambda_L = 0$
- **outside the DV**
 $b_L = -0.010$
 $b_R = +0.030$
 $\lambda_L = 0$
- **$t\bar{b}$ & $b\bar{b}$ in the DV**
 $b_L = +0.009$
 $b_R = +0.030$
 $\lambda_L = +0.006$
- **all in the DV**
 $b_L = +0.0098$
 $b_R = +0.0034$
 $\lambda_L = +0.006$



HIDING THE PEAK

$$M_V = 1 \text{ TeV}, \ g'' = 20, \ p = 0, \ \lambda_R = 0$$



- **no direct cplng**
 $b_L = 0$
 $b_R = 0$
 $\lambda_L = 0$
- **outside the DV**
 $b_L = -0.010$
 $b_R = +0.030$
 $\lambda_L = 0$
- **$t\bar{b}$ & $b\bar{b}$ in the DV**
 $b_L = +0.009$
 $b_R = +0.030$
 $\lambda_L = +0.006$
- **all in the DV**
 $b_L = +0.0098$
 $b_R = +0.0034$
 $\lambda_L = +0.006$



$pp \rightarrow t\bar{b} + X$ @ LHC

$\sqrt{s} = 14 \text{ TeV}, \Gamma_{V^\pm} = 50 \text{ GeV}$


$|y(t)|, |y(\bar{b})| < 2.5, \quad p_T(t), p_T(\bar{b}) > 200 \text{ GeV}$

PSP	g''	p	b_L	b_R	R
A	20	0	-0.15	N/A	11.6
B	20	0	+0.16	N/A	12.1
C	20	1	-0.11	-0.11	10.7
D	30	0	-0.11	N/A	6.5
E	20	1	-0.11	+0.11	10.6

$R = \frac{N_{tBESS} - N_{SM}}{\sqrt{N_{SM}}}$

$N = \sigma L \epsilon_1 \epsilon_b^2$

$L = 100 \text{ fb}^{-1}, \epsilon_1 = 0.216,$
 $\epsilon_b = 0.4$



SUMMARY

- effective description of strong ESB new physics
- top-BESS — modification of BESS, special role of top quark
 - ◊ *new $SU(2)$ resonance triplet*
 - ◊ *direct coupling to top and bottom*
 - ◊ *λ -terms*
- low-E limits on the fermion parameters relaxed
- the Death Valley effect
- LHC: $pp \rightarrow t\bar{b}X$

BACKUP


BACKUP SLIDES

NEW RESONANCE MASSES

- mass of the vector resonance:

$$M_V = \frac{\sqrt{\alpha} g'' v}{2}$$

- EW gauge bosons \rightarrow mixing \rightarrow mass splitting



LOW-ENERGY LIMIT FOR g''

THEORY:


$$\epsilon_3 \approx \epsilon_3^{tree} + \underbrace{\left(5.25 + 0.54 \ln \frac{M_H}{M_Z} \right) \times 10^{-3}}_{\epsilon_3^{SM \ loop}}$$

M_H (TeV)	0.3	1	2	$\epsilon_3^{exp} \times 10^3$
$\epsilon_3^{SM \ loop} \times 10^3$	5.89	6.54	6.92	5.34 ± 0.94

$$\Rightarrow \epsilon_3^{tree} < 0$$

$$\epsilon_3^{tree}(BESS) = -\frac{b_L}{2} + \left(\frac{g}{g''} \right)^2$$

$$\epsilon_3^{tree}(tBESS) = \left(\frac{g}{g''} \right)^2$$



NP FERMION INVARIANTS

$$I_c^L = i\bar{\psi}_L(\partial + \mathbf{W} + \mathbf{B})\psi_L$$

$$I_c^R = i\bar{\psi}_R(\partial + \mathbf{B})\psi_R$$

$$I_b^h = i\bar{\chi}_h(\partial + \mathbf{V} + ig'\mathcal{B}(B-L)/2)\chi_h$$

$$\begin{aligned} I_\lambda^h &= i\bar{\chi}_h \bar{\psi}^\perp \chi_h \\ &= i\bar{\chi}_h [\psi^\perp + (\xi_L^\dagger \mathbf{W} \xi_L - \xi_R^\dagger \mathcal{B}^{R3} \xi_R)] \chi_h \end{aligned}$$

$$\chi_h = \xi_h^\dagger \psi_h, \quad h = L, R$$