





Chris Kouvaris





Particle Physics & Origin of Mass

Planck 2012

Wednesday, May 30, 2012



The Missing Mass of the Universe

A Mystery for 80 years!



Zwicky 1933: Virial Theorem on Coma cluster



Vera Rubin 70's Rotation curves of Andromeda are not falling according to Newton's law!



Microwave Background Radiation





Bullet Cluster



What is the nature of Dark Matter?

WIMP-nucleus cross section:

- •Spin-Independent
- •Spin-dependent
- Inelastic cross section

WIMP-WIMP cross section:Self-Interacting dark matter

WIMP-WIMP annihilation:

- •Thermally produced WIMPs
- •Nonthermally, asymmetric dark matter

Decaying WIMPs: possible explanation of PAMELA results



Astrophysical Observations



Astrophysical Observations

WIMP annihilation and Cooling of Stars

WIMP annihilation as a heating mechanism for
neutron stars (CK '07, CK Tinyakov '10, Lavallaz Fairbairn '10)
white dwarfs (Bertone Fairbairn '07, McCullough '10)



Astrophysical Observations

WIMP annihilation and Cooling of Stars

WIMP annihilation as a heating mechanism for
neutron stars (CK '07, CK Tinyakov '10, Lavallaz Fairbairn '10)
white dwarfs (Bertone Fairbairn '07, McCullough '10)

WIMP collapse to a Black Hole

WIMPs can be trapped inside stars and later collapse forming a black hole that destroys the star (Goldman Nussinov '89, CK Tinyakov '10, '11, McDermott Yu Zurek '11, CK'11, Guver Erkoca Reno Sarcevic '12, Fan Yang Chang '12)



WIMP capture in Stars

<u>Condition:</u> The energy loss in the collision should be larger than the asymptotic kinetic energy of the WIMP far out of the star.



Example: Sun

WIMP mean free path inside the sun $\xi \approx \frac{1}{n\sigma}$, $n \approx \frac{M_{solar}}{(4/3)\pi R_{solar}^3 m_n} \approx 8.10^{23}$ particles/cm³

Even if current $\sigma < 10^{-7}$ limit of CDMS

$$^{-41}cm^2$$
, $\xi \approx 10^{17}cm$, $\frac{R_{solar}}{\varepsilon} \approx 10^{17}cm$

Only one out of a million WIMPs scatters!



For a typical neutron star $M_{NS} \approx 1.4 M_{solar}, R \approx 10 km$

$$\sigma > \sigma_{critical} \approx 5 \cdot 10^{-46} cm^2$$
 CK'07

For cross section larger than the critical one, every WIMP passing through the neutron star will be on average interact inside the star.







Wednesday, May 30, 2012













Wednesday, May 30, 2012





































WIMP capture in Stars

$$F = \frac{8}{3}\pi^2 \frac{\rho_{\rm dm}}{m} \left(\frac{3}{2\pi v^2}\right)^{3/2} \frac{GMR}{1 - \frac{2GM}{R}} v^2 (1 - e^{-3E_0/v^2}) f \quad \text{CK'07}$$

For typical NS $F = 1.25 \times 10^{24} \text{s}^{-1} \left(\frac{\rho_{\text{dm}}}{\text{GeV/cm}^3} \right) \left(\frac{100 \text{GeV}}{m} \right) f$

Thermalization

$$t_{\rm th} = 0.2 {
m yr} \left(\frac{m}{{
m TeV}}\right)^2 \left(\frac{\sigma}{10^{-43} {
m cm}^2}\right)^{-1} \left(\frac{T}{10^5 {
m K}}\right)^{-1}$$

Goldman Nussinov'89

$$r_{\rm th} = \left(\frac{9T}{8\pi G\rho_c m}\right)^{1/2} \simeq 22 {\rm cm} \left(\frac{T}{10^5 {\rm K}}\right)^{1/2} \left(\frac{100 {\rm GeV}}{m}\right)^{1/2}$$

Wednesday, May 30, 2012

Danish Institute for Advanced Study

Cooling of Neutron Stars

$$\frac{dT}{dt} = \frac{-L_{\nu} - L_{\gamma} + L_{\rm dm}}{Vc_V} = \frac{V(-\epsilon_{\nu} - \epsilon_{\gamma} + \epsilon_{\rm dm})}{Vc_V} = \frac{-\epsilon_{\nu} - \epsilon_{\gamma} + \epsilon_{\rm dm}}{c_V}$$



CK'07

Wednesday, May 30, 2012

Danish Institute for Advanced Study

Cooling of Neutron Stars

$$\frac{dT}{dt} = \frac{-L_{\nu} - L_{\gamma} + L_{\rm dm}}{Vc_V} = \frac{V(-\epsilon_{\nu} - \epsilon_{\gamma} + \epsilon_{\rm dm})}{Vc_V} = \frac{-\epsilon_{\nu} - \epsilon_{\gamma} + \epsilon_{\rm dm}}{c_V}$$



CK'07

Cooling of Neutron Stars

$$\frac{dT}{dt} = \frac{-L_{\nu} - L_{\gamma} + L_{\rm dm}}{Vc_V} = \frac{V(-\epsilon_{\nu} - \epsilon_{\gamma} + \epsilon_{\rm dm})}{Vc_V} = \frac{-\epsilon_{\nu} - \epsilon_{\gamma} + \epsilon_{\rm dm}}{c_V}$$



Cooling of Neutron Stars

$$\frac{dT}{dt} = \frac{-L_{\nu} - L_{\gamma} + L_{\rm dm}}{Vc_V} = \frac{V(-\epsilon_{\nu} - \epsilon_{\gamma} + \epsilon_{\rm dm})}{Vc_V} = \frac{-\epsilon_{\nu} - \epsilon_{\gamma} + \epsilon_{\rm dm}}{c_V}$$





Cooling of Neutron Stars

Galactic Center



FIG. 3: The surface temperature of a typical old neutron star in units of K as a function of the distance of the star from the galactic center in pc, with the dark matter annihilation taken into account. The three curves correspond to three different dark matter profiles: NFW (thin solid line), Einasto (thick solid line), and Burkert (dashed line).

Globular Cluster



FIG. 5: The surface temperature of a typical old neutron star in units of K as a function of the distance in pc for a NFW profile of the globular cluster M4.

$$\rho_{\rm NFW} = \frac{\rho_s}{\frac{r}{r_s}(1+\frac{r}{r_s})^2} \qquad \rho_{\rm Ein} = \rho_s \exp\left[-\frac{2}{\alpha} \left[\left(\frac{r}{r_s}\right)^{\alpha} - 1\right]\right] \qquad \rho_{\rm Bur} = \frac{\rho_s}{\left(1+\frac{r}{r_s}\right) \left[1+\left(\frac{r}{r_s}\right)^2\right]}$$

Nearby old neutron stars

J0437-4715 temperature ~10^5 K J2124-3358 temperature ~10^5 K 130-140 pc away CK, Tinyakov '10

Cooling of Neutron Stars



Old neutron stars in Globular Clusters

X7 in 47 Tuc 1620-26 in M4

both have temperatures roughly 10^6 K







$$\frac{GNm^2}{r} \simeq \frac{\hbar}{r} \longrightarrow M_{crit} = \frac{2M_{\rm Pl}^2}{\pi m} \sqrt{1 + \frac{M_{\rm Pl}^2}{4\sqrt{\pi}m} \sigma^{1/2}} \qquad \sigma = \lambda^2 / (64\pi m^2)$$



$$\frac{GNm^2}{r} \simeq \frac{\hbar}{r} \longrightarrow M_{crit} = \frac{2M_{\rm Pl}^2}{\pi m} \sqrt{1 + \frac{M_{\rm Pl}^2}{4\sqrt{\pi m}} \sigma^{1/2}} \qquad \sigma = \lambda^2/(64\pi m^2)$$
repulsive interactions

Bosonic Asymmetric Dark Matter

$$\frac{GNm^2}{r} \simeq \frac{\hbar}{r} \longrightarrow M_{crit} = \frac{2M_{\rm Pl}^2}{\pi m} \sqrt{1 + \frac{M_{\rm Pl}^2}{4\sqrt{\pi m}} \sigma^{1/2}} = \frac{\lambda^2}{(64\pi m^2)}$$
BEC accelerates collapse
$$T_c = \left(\frac{n}{\zeta(3/2)}\right)^{2/3} \frac{2\pi\hbar^2}{mk_B} \approx 3.31 \frac{\hbar^2 n^{2/3}}{mk_B} \qquad N_{\rm BEC} \simeq 2 \times 10^{36}$$

$$r_{\rm th} \simeq 2 \,\mathrm{m} \left(\frac{T_c}{10^5 \mathrm{K}}\right)^{1/2} \left(\frac{m}{\mathrm{GeV}}\right)^{-1/2} \longrightarrow r_c = \left(\frac{8\pi}{3}G\rho_c m^2\right)^{-1/4} \simeq 1.6 \times 10^{-4} \left(\frac{\mathrm{GeV}}{m}\right)^{1/2} \mathrm{cm}$$

Bosonic Asymmetric Dark Matter





Bosonic Asymmetric Dark Matter



CK, Tinyakov PRL '11

Bosonic Asymmetric Dark Matter



 $D|\Lambda S$ Danish Institute for Advanced Study

Bosonic Asymmetric Dark Matter

 $\rho=10^4 \text{GeV/cm}^3$

excluded by neutron stars

10³

 $\rho=~10^3 \text{GeV/cm}^{-3}$

10⁶

Hawking

radiation

prevents

BH growth

m, GeV



Bosonic Asymmetric Dark Matter



If WIMP is a composite of fermions

$$\Lambda_{crit} = m^{1/3} M_{\rm Pl}^{2/3} \left(1 + \frac{\lambda m_{pl}^2}{32\pi m^2} \right)^{-1/3}$$

If WIMP is a composite of fermions above that scale, the bosonic constraints still hold



Self-Interacting Dark Matter

"Chandrashekhar Limit for WIMPs"

$$\frac{GNm^2}{r} > k_F = \left(\frac{3\pi^2 N}{V}\right) = \left(\frac{9\pi}{4}\right)^{1/3} \frac{N^{1/3}}{r} \qquad N = 10^{57} / m^{3}!!!$$

Yukawa-type WIMP self-interactions can explain the flatness of dwarf galaxies Spergel-Steinhardt '99, Loeb-Weiner '11

$$\alpha \phi \bar{\psi} \psi$$
 $V(r) = -\alpha \exp[-\mu r]/r$

Yukawa self-interactions can alleviate the effect of the Fermi pressure, leading to a gravitational collapse with dramatically lower amount of captured WIMPs



Self-Interacting Dark Matter



Wednesday, May 30, 2012



Wednesday, May 30, 2012



A regular star accumulates WIMPs with spin-dependent WIMP-nucleon interactions and collapses to a white dwarf after the hydrogen and helium burning stages



A regular star accumulates WIMPs with spin-dependent WIMP-nucleon interactions and collapses to a white dwarf after the hydrogen and helium burning stages

The WIMP population is inherited by the white dwarf and gets thermalized inside it due to the presence of CI3-WIMP spin-dependent interactions



A regular star accumulates WIMPs with spin-dependent WIMP-nucleon interactions and collapses to a white dwarf after the hydrogen and helium burning stages

The WIMP population is inherited by the white dwarf and gets thermalized inside it due to the presence of CI3-WIMP spin-dependent interactions



Formation of a Black Hole

The Dark Side of the Stars

Compact stars can reveal a lot of information about the nature of DM putting constraints on its properties complementary to direct searches.

• Observation of cold neutron stars can exclude thermally produced dark matter.

• Asymmetric dark matter:

I. keV to ~I6GeV bosonic dark matter is excluded.
2.Part of fermionic WIMP self-interactions excluded.
3.Constraints on WIMP-nucleon spin-dependent interactions.