Fermiophobic Higgs boson and supersymmetry

Antonio Racioppi

NICPB, Tallinn, Estonia

Warsaw, May 30th, 2012

based on 1204.0080 in collaboration with

- E. Gabrielli (NICPB),
- K. Kannike (NICPB & SNS and INFN, Pisa, Italy),
- B. Mele (INFN, Rome, Italy),
- M. Raidal (NICPB & Institute of Physics, Tartu, Estonia & CERN)

イロン イヨン イヨン イヨン

Motivation MSSM & Higgs mass loop correction The failure of the MSSM

Motivation

Why fermiophobia?

- ► Th: Chiral Symmetry Breaking (m_f) and EW symmetry breaking (M_W, M_Z) may have a different origin.
 ⟨h⟩ could be only responsible of M_W and M_Z but not of m_f
- **Exp 1**: not (yet) any strong evidence supporting tree-level y_f
- ► **Exp 2**: local excess in $\gamma\gamma$ at LHC $\stackrel{?}{\Rightarrow}$ FP *h*, $M_h \simeq 125$ GeV CMS-PAS-HIG-12-002,008; ATLAS-CONF-2012-013

Why supersymmetry?

Only NON susy fermiophobic Higgs models in the past

イロト イポト イヨト イヨト

Motivation MSSM & Higgs mass loop correction The failure of the MSSM

Motivation

Pros:

- sfermions can be naturally in the 2-3 TeV range
- the usual flavor and CP problems are relaxed
- ▶ no more constraints from $b \rightarrow s\gamma$ and $B_s \rightarrow \mu\mu$ ⇒ H^{\pm} can be light
- relax constraints on DM

Cons:

► No mechanism for generating SM m_f (added by hand) ⇒ it is only an effective theory

イロト イポト イヨト イヨト

Motivation MSSM & Higgs mass loop correction The failure of the MSSM

MSSM & Higgs mass loop correction

In the MSSM $M_h^{\text{tree}} < M_Z$. But

$$\Delta M_h^2 = 3y_t^4 \frac{v^2 \sin^4 \beta}{8\pi^2} \left[\log \frac{M_S^2}{m_t^2} + \frac{X_t^2}{M_S^2} \left(1 - \frac{X_t^2}{12M_S^2} \right) \right]$$

- *M_S* is the average stop mass
- X_t is the stop mass mixing parameter, $X_t = \frac{a_t}{v_t} \mu \cot \beta$
- a_t is the trilinear coupling of the soft term $a_t \tilde{Q} H_u \tilde{u}^c$

・ロト ・回ト ・ヨト ・ヨト

Motivation MSSM & Higgs mass loop correction The failure of the MSSM

The failure of the MSSM

Fermiophobic limit: $y_t \rightarrow 0$

$$\Delta M_h^2 = 3 y_t^4 \frac{v^2 \sin^4 \beta}{8\pi^2} \left[\log \frac{M_S^2}{m_t^2} + \frac{X_t^2}{M_S^2} \left(1 - \frac{X_t^2}{12M_S^2} \right) \right]$$

- *M_S* is the average stop mass
- X_t is the stop mass mixing parameter, $X_t = \frac{a_t}{v_t} \mu \cot \beta$
- a_t is the trilinear coupling of the soft term $a_t \tilde{Q} H_u \tilde{u}^c$

イロト イポト イヨト イヨト

Motivation MSSM & Higgs mass loop correction The failure of the MSSM

The failure of the MSSM

Fermiophobic limit: $y_t \rightarrow 0$

$$\Delta M_h^2 = -\frac{3v^2 \sin^4 \beta}{8\pi^2} \frac{a_t^4}{12M_S^4} < 0$$

- ▶ so we cannot get $M_h > M_Z$ and even more so $M_h \simeq 125$ GeV
- $M_h < M_Z$ is already ruled out
- ► Fermiophobic Higgs if SUSY, cannot be just MSSM-like

Setup Scalar potential and masses Neutralinos & Charginos

Setup

 Z_3 symmetric FP NMSSM superpotential

$$\mathcal{W} = \lambda SH_uH_d + \frac{k}{3}S^3$$

and S, H_u, H_d soft terms

$$\mathcal{L}_{\text{soft}}^{S,H} = -\left(m_{h_u}^2 h_u^{\dagger} h_u + m_{h_d}^2 h_d^{\dagger} h_d + m_s^2 s^{\dagger} s\right) - \left(a_\lambda s h_u h_d + \frac{1}{3}a_k s^3 + h.c.\right)$$

where

- s is the scalar component of S
- Z_3 is also used in order to get fermiophobia

►
$$X_{H_u} = X_{H_d} = X_S = 1$$
 and $X_f = 0 \Rightarrow y_f, a_f = 0$
But also other possible configurations (see 1204.0080)

Setup Scalar potential and masses Neutralinos & Charginos

Scalar potential

$$\begin{split} V &= \left(m_{h_u}^2 + |\lambda s|^2\right) \left(\left|h_u^0\right|^2 + \left|h_u^+\right|^2\right) + \left(m_{h_d}^2 + |\lambda s|^2\right) \left(\left|h_d^0\right|^2 + \left|h_d^-\right|^2\right) \\ &+ m_s^2 |s|^2 + \left(a_\lambda \left(h_u^+ h_d^- - h_u^0 h_d^0\right) s + \frac{1}{3} a_k s^3 + \text{h.c.}\right) \\ &+ \left|\lambda \left(h_u^+ h_d^- - h_u^0 h_d^0\right) + k s^2\right|^2 \\ &+ \frac{g_1^2 + g_2^2}{8} \left(\left|h_u^0\right|^2 + \left|h_u^+\right|^2 - \left|h_d^0\right|^2 - \left|h_d^-\right|^2\right)^2 + \frac{g_2^2}{2} \left|h_u^+ h_d^{0*} + h_u^0 h_d^{-*}\right|^2 \end{split}$$

Parametrization:

$$h_{d}^{0} = \frac{1}{\sqrt{2}} \left(v_{d} + h_{dR}^{0} + ih_{dl}^{0} \right) \qquad v^{2} = v_{u}^{2} + v_{d}^{2}$$

$$h_{u}^{0} = \frac{1}{\sqrt{2}} \left(v_{u} + h_{uR}^{0} + ih_{ul}^{0} \right) \qquad \tan \beta = \frac{v_{u}}{v_{d}}$$

$$s = \frac{1}{\sqrt{2}} \left(v_{S} + s_{R} + is_{l} \right)$$

æ

Setup Scalar potential and masses Neutralinos & Charginos

CP even scalars

$$M_{5}^{2} = \begin{pmatrix} M_{5,11}^{2} & M_{5,12}^{2} & M_{5,13}^{2} \\ \dots & M_{5,22}^{2} & M_{5,23}^{2} \\ \dots & \dots & M_{5,33}^{2} \end{pmatrix} \text{ in the basis } (h_{dR}^{0}, h_{uR}^{0}, s_{R})$$

$$M_{5,13}^2 = vv_S \left(\lambda^2 \cos\beta - k\lambda \sin\beta\right) - \frac{a_\lambda v \sin\beta}{\sqrt{2}}$$
$$M_{5,23}^2 = vv_S \left(\lambda^2 \sin\beta - k\lambda \cos\beta\right) - \frac{a_\lambda v \cos\beta}{\sqrt{2}}$$

There is a choice that allows no mixing between s_R and $h_{uR,dR}^0$,

$$egin{array}{l} a_\lambda = 0 \ k = \lambda \ ext{tan} \ eta = 1 \end{array}$$

N.B. tan $\beta = 1$ is allowed in this model because no constraints occur from the Yukawa sector. Therefore this choice is the most natural one $\beta + 2\beta + 2\beta + 2\beta$.

Setup Scalar potential and masses Neutralinos & Charginos

CP even scalars

Then the minimization equations read

$$\begin{array}{rcl} \left(4m_{h_d}^2 + \lambda^2 v^2\right) &=& 0\\ \left(4m_{h_u}^2 + \lambda^2 v^2\right) &=& 0\\ \\ \frac{a_k v_S^2}{\sqrt{2}} + m_S^2 v_S + \lambda^2 v_S^3 &=& 0 \end{array}$$

leading to the CP-even neutral Higgs boson mass matrix

$$M_{5}^{2} = \begin{pmatrix} \frac{1}{2} \left(M_{Z}^{2} + v_{5}^{2} \lambda^{2} \right) & \frac{1}{2} \left((v - v_{5})(v + v_{5})\lambda^{2} - M_{Z}^{2} \right) & 0 \\ \dots & \frac{1}{2} \left(M_{Z}^{2} + v_{5}^{2} \lambda^{2} \right) & 0 \\ \dots & \dots & 2v_{5}^{2} \lambda^{2} + \frac{a_{k}v_{5}}{\sqrt{2}} \end{pmatrix}$$

where $M_Z^2 = rac{1}{4} \left(g_1^2 + g_2^2
ight) v^2.$

・ロン ・回と ・ヨン・

Setup Scalar potential and masses Neutralinos & Charginos

CP even scalars

The corresponding eigenvectors and eigenvalues

$$h = \frac{1}{\sqrt{2}} (h_{dR}^{0} + h_{uR}^{0}) \qquad M_{h}^{2} = \frac{(\lambda v)^{2}}{2} \simeq (125 \text{GeV})^{2}$$
$$H = \frac{1}{\sqrt{2}} (h_{dR}^{0} - h_{uR}^{0}) \qquad M_{H}^{2} = (\lambda v_{S})^{2} + M_{Z}^{2} - M_{h}^{2}$$
$$M_{s_{R}}^{2} = \frac{a_{k} v_{S}}{\sqrt{2}} + 2(\lambda v_{S})^{2}$$

 $\begin{array}{c} \text{MSSM notation} \rightarrow \alpha = -\pi/4 \\ \downarrow \\ \beta = \pi/4 \Rightarrow \text{ no direct tree level couplings } HWW \text{ and } HZZ \\ \downarrow \\ \text{OK with LHC seeing no other "resonance" but } \simeq 125 \text{GeV.} \\ \hline \end{array}$

Setup Scalar potential and masses Neutralinos & Charginos

CP odd scalars

$$\mathcal{M'}_{P}^{2} = \begin{pmatrix} \frac{v_{S}^{2}\lambda^{2}}{2} & \frac{v_{S}^{2}\lambda^{2}}{2} & -\frac{v_{VS}\lambda^{2}}{\sqrt{2}} \\ \frac{v_{S}^{2}\lambda^{2}}{2} & \frac{v_{S}^{2}\lambda^{2}}{2} & -\frac{v_{VS}\lambda^{2}}{\sqrt{2}} \\ -\frac{v_{VS}\lambda^{2}}{\sqrt{2}} & -\frac{v_{VS}\lambda^{2}}{\sqrt{2}} & v^{2}\lambda^{2} - \frac{3a_{k}v_{S}}{\sqrt{2}} \end{pmatrix}$$

in the basis
$$(h_{dI}^0, h_{uI}^0, s_I)$$

Э

and the corresponding eigenvalues

$$M_{G^0}^2 = 0$$

$$M_{A_{1,2}^0}^2 = \frac{1}{4} \left(2\lambda^2 \left(v^2 + v_5^2 \right) - 3\sqrt{2}a_k v_5 \mp \sqrt{\Delta} \right)$$

where

$$\Delta = \left(3\sqrt{2}a_kv_S - 2\lambda^2\left(v^2 + v_S^2\right)\right)^2 + 24\sqrt{2}a_k\lambda^2v_S^3$$

Setup Scalar potential and masses Neutralinos & Charginos

Charged scalars

$$\begin{split} M'_{\pm}^2 &= \left(M_W^2 + \frac{1}{2} \lambda^2 \left(2v_5^2 - v^2 \right) \right) \left(\begin{array}{cc} 1 & 1 \\ 1 & 1 \end{array} \right) & \text{ in the basis } (h_u^+, h_d^{-*}) \end{split}$$
where $M_W^2 &= \frac{1}{4} g_2^2 v^2.$

It contains one massless Goldstone mode G^{\pm} , and

イロン イボン イヨン イヨン 三日

Setup Scalar potential and masses Neutralinos & Charginos

Neutralinos & Charginos

$$M_0 = \begin{pmatrix} M_1 & 0 & -M_Z \sin \theta_W & M_Z \sin \theta_W & 0\\ \dots & M_2 & M_Z \cos \theta_W & -M_Z \cos \theta_W & 0\\ \dots & \dots & 0 & -\frac{1}{\sqrt{2}} \lambda v_S & -\frac{\lambda v}{2}\\ \dots & \dots & 0 & -\frac{\lambda v}{\sqrt{2}} \\ \dots & \dots & \dots & 0 & -\frac{\lambda v}{2} \\ \dots & \dots & \dots & \sqrt{2} v_S \end{pmatrix}$$

in the basis $\psi^0 = (\lambda_1, \lambda_2^3, \tilde{h}_d^0, \tilde{h}_u^0, \tilde{s})$

$$M_{\tilde{C}} = \begin{pmatrix} \mathbf{0} & \mathbf{X}^{T} \\ \mathbf{X} & \mathbf{0} \end{pmatrix} \text{ in the basis } \psi^{\pm} = (\tilde{W}^{+}, \tilde{H}_{u}^{+}, \tilde{W}^{-}, \tilde{H}_{d}^{-})$$
$$\mathbf{X} = \begin{pmatrix} M_{2} & M_{W} \\ M_{W} & \frac{1}{\sqrt{2}}\lambda v_{S} \end{pmatrix}$$

Strategy Plots & Results

Strategy

►



relevant free parameters:

 $M_1 > 0$, $M_2 > 0$, $|\lambda|$, $|\mu| \equiv |\lambda v_S| / \sqrt{2}$, sign (μ)

- 4 回 2 4 三 2 4 三 2 4

æ

Strategy Plots & Results

Strategy



► relevant free parameters: $M_1 > 0$, $M_2 > 0$, $|\lambda|$, $|\mu| \equiv |\lambda v_S| / \sqrt{2}$, sign(μ)

•
$$M_h = \frac{|\lambda|v}{\sqrt{2}} = 125$$
 GeV, $M_1 = 100$ GeV

- 4 回 2 4 三 2 4 三 2 4

æ

Strategy Plots & Results

Strategy



▶ relevant free parameters: $M_1 > 0$, $M_2 > 0$, $|\lambda|$, $|\mu| \equiv |\lambda v_S| / \sqrt{2}$, sign(μ)

•
$$M_h = \frac{|\lambda|v}{\sqrt{2}} = 125$$
 GeV, $M_1 = 100$ GeV

$$\blacktriangleright (M_{H^{\pm}}, M_{\chi_L^+}) \rightarrow (|\mu|, M_2)$$

イロン イヨン イヨン イヨン

Strategy Plots & Results

Strategy



► relevant free parameters: $M_1 > 0$, $M_2 > 0$, $|\lambda|$, $|\mu| \equiv |\lambda v_S| / \sqrt{2}$, sign(μ)

•
$$M_h = rac{|\lambda| v}{\sqrt{2}} = 125$$
 GeV, $M_1 = 100$ GeV

$$\blacktriangleright (M_{H^{\pm}}, M_{\chi_L^+}) \to (|\mu|, M_2)$$

$$M_{\chi_{L}^{0}} > M_{h}/2 \Rightarrow h \nleftrightarrow \chi_{i}\chi_{j} \Rightarrow R - \text{parity} \Rightarrow h \nleftrightarrow \chi_{i}^{*}\chi_{j}, \chi_{i}^{*}\chi_{j}$$

Strategy Plots & Results

Signal rates @ LHC7



- 1. The lines above (below) the FP SM prediction for $h \rightarrow \gamma \gamma$ correspond to positive (negative) values the μ . For $h \rightarrow Z \gamma$ is the opposite.
- 2. For fixed $M_{\chi_L^+}$ two non-degenerates values of $M_{\chi_H^+}$ are possible. So there are always two solutions for the one-loop SUSY contribution.

A⊒ ▶ ∢ ∃

Strategy Plots & Results

Signal rates @ LHC7



- 3. As in the MSSM, the dominant SUSY contribution comes from the χ^{\pm}
- 4. The absence of points in the half-plane above (below) the FP(SM) line for $M_{H^+} = 200$ GeV in the case of $h \rightarrow \gamma\gamma$ ($h \rightarrow Z\gamma$), is due to the constraint $M_{\chi_1^0} > M_h/2$ and depends on our choice for $M_1 = 100$ GeV.

Strategy Plots & Results

Signal rates @ LHC7



5. The present fits (1203.4254) indicate that the LHC observes fewer $\gamma\gamma$ events than predicted by the pure FP SM. FP NMSSM signal rates can be smaller than the FP SM predictions, so FP NMSSM allows a better fit to LHC data.

イロト イポト イヨト イヨト

э

Conclusions

Results:

- Fermiophobic Higgs boson in MSSM is ruled out
- FP NMSSM is viable and Z_3 could explain fermiophobia
- 1-loop Higgs boson branching fractions and production rates in $\gamma\gamma$ and $Z\gamma$ can be sizably modified with respect to the FP SM, allowing a better fit to present collider data

Future:

- Admitting the s_R − h⁰_{uR,dR} mixing the Higgs coupling to weak gauge boson WW and ZZ can be modified, and a suppression of the rates for h → WW* and h → ZZ* with respect to the pure FP model expectations can be achieved.
- Most of the previous analyses on SUSY particle searches should be revised in the light of the fact that the large top-Yukawa coupling is absent or strongly suppressed.

Thank you!

Backup slides

Antonio Racioppi Fermiophobic Higgs boson and supersymmetry

< □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □

Tevatron



イロン イヨン イヨン イヨン

Э

Fermion mass

The obvious question in any FP Higgs boson scenario is what is the alternative mechanism for generating the observed fermion masses. Because the top quark mass is so large, it cannot be generated radiatively. The most plausible scenario for generating such large fermion masses is strong dynamics above the electroweak scale. In such a scenario both the composite Higgs boson fermiophobia and fermion masses might originate from the same new physics. See: hep-ph/0703164, 1002.1011, 1012.1562, 1110.1613

A generic prediction of strong electroweak symmetry breaking scenarios, including composite Higgs models, is the appearance of new resonances at 2–3 TeV. We assume that such or any other new physics scenario above the electroweak scale generates the top quark mass.

イロト イポト イヨト イヨト

XENON100

In the MSSM the spin-independent dark matter scattering off nuclei is dominated by tree level Higgs boson exchange. Now this process is suppressed. The dominant dark matter scattering process is through WW exchange at one loop level. Scattering due to W-loops is too weak to be observed in the present stage of XENON100. As the SUSY scale could now be large, higgsino or wino relic abundance would become a natural explanation to the dark matter of the Universe.

- 4 回 ト 4 ヨ ト 4 ヨ ト

Z_3 charge assignment

• $\mu H_u H_d$, S, S², $y_u Q H_u u^c$, $y_d H_d Q d^c$, $y_e H_d L e^c$ forbidden by:

 $\begin{array}{l} X_Q + X_{H_u} + X_{u^c} \neq 0 \mod N, \\ X_Q + X_{H_d} + X_{d^c} \neq 0 \mod N, \\ X_L + X_{H_d} + X_{e^c} \neq 0 \mod N, \\ X_{H_u} + X_{H_d} \neq 0 \mod N, \\ X_S \neq 0 \mod N, \\ 2X_S \neq 0 \mod N, \end{array}$

• λSH_uH_d , $\frac{k}{3}S^3$ allowed by

 $3X_S = 0 \mod N,$ $X_S + X_{H_u} + X_{H_d} = 0 \mod N$

Easiest: $X_{H_u} = X_{H_d} = X_S = 1$ and $X_{\text{fermion}} = 0 \Rightarrow \text{Yukawas via } \frac{H_u H_d}{\Lambda^2} Q H_d d^c \dots$ Other: $X_{H_u} = X_{H_d} = X_S = 1$ and $X_{\text{fermion}} = 2 \Rightarrow \text{Yukawas via } \frac{S}{\Lambda} Q H_d d^c \dots$

Scalar potential minimization

Parametrization:

$$\begin{split} h_{d}^{0} &= \frac{1}{\sqrt{2}} \left(v_{d} + h_{dR}^{0} + i h_{dI}^{0} \right) & v^{2} &= v_{u}^{2} + v_{d}^{2} \\ h_{u}^{0} &= \frac{1}{\sqrt{2}} \left(v_{u} + h_{uR}^{0} + i h_{uI}^{0} \right) & \tan \beta &= \frac{v_{u}}{v_{d}} \\ s &= \frac{1}{\sqrt{2}} \left(v_{S} + s_{R} + i s_{I} \right) \end{split}$$

Minima equations:

$$v\left(-4v_{S}\sin\beta\left(\sqrt{2}a_{\lambda}+k\lambda v_{S}\right)+\cos\beta\left(v^{2}\cos^{3}\beta\left(g_{1}^{2}+g_{2}^{2}\right)+8m_{h_{d}}^{2}+4\lambda^{2}v_{S}^{2}\right)\right)$$
$$-v^{2}\sin^{2}\beta\left(g_{1}^{2}+g_{2}^{2}-4\lambda^{2}\right)\right)=0$$
$$v\left(-4v_{S}\cos\beta\left(\sqrt{2}a_{\lambda}+k\lambda v_{S}\right)+\sin\beta\left(v^{2}\sin^{2}\beta\left(g_{1}^{2}+g_{2}^{2}\right)+8m_{h_{u}}^{2}+4\lambda^{2}v_{S}^{2}\right)\right)$$
$$-v^{2}\sin\beta\cos^{2}\beta\left(g_{1}^{2}+g_{2}^{2}-4\lambda^{2}\right)\right)=0$$
$$v_{S}\left(\sqrt{2}a_{k}v_{S}+2k^{2}v_{S}^{2}+\lambda^{2}v^{2}+2m_{S}^{2}\right)-\frac{1}{4}v^{2}\sin2\beta\left(\sqrt{2}a_{\lambda}+2k\lambda v_{S}\right)=0$$

CP even scalars

$$M_{5}^{2} = \begin{pmatrix} M_{5,11}^{2} & M_{5,12}^{2} & M_{5,13}^{2} \\ \dots & M_{5,22}^{2} & M_{5,23}^{2} \\ \dots & \dots & M_{5,33}^{2} \end{pmatrix} \text{ in the basis } (h_{dR}^{0}, h_{uR}^{0}, s_{R})$$

where

$$\begin{split} M_{S,11}^2 &= m_{h_d}^2 + \frac{v_S^2 \lambda^2}{2} + \frac{1}{8} v^2 \left(g_1^2 + g_2^2 + 2\lambda^2 + 2 \left(g_1^2 + g_2^2 - \lambda^2 \right) \cos 2\beta \right) \\ M_{S,22}^2 &= m_{h_u}^2 + \frac{v_S^2 \lambda^2}{2} + \frac{1}{8} v^2 \left(g_1^2 + g_2^2 + 2\lambda^2 - 2 \left(g_1^2 + g_2^2 - \lambda^2 \right) \cos 2\beta \right) \\ M_{S,33}^2 &= m_S^2 + 3k^2 v_S^2 + \sqrt{2} a_k v_S + v^2 \left(\frac{\lambda^2}{2} - k\lambda \cos \beta \sin \beta \right) \\ M_{S,12}^2 &= \frac{1}{8} \left(-g_1^2 - g_2^2 + 4\lambda^2 \right) \sin 2\beta v^2 - \frac{1}{2} k v_S^2 \lambda - \frac{a_\lambda v_S}{\sqrt{2}} \\ M_{S,13}^2 &= v v_S \left(\lambda^2 \cos \beta - k\lambda \sin \beta \right) - \frac{a_\lambda v \sin \beta}{\sqrt{2}} \\ M_{S,23}^2 &= v v_S \left(\lambda^2 \sin \beta - k\lambda \cos \beta \right) - \frac{a_\lambda v \cos \beta}{\sqrt{2}} \\ \end{split}$$

Antonio Racioppi

CP even scalars eigenstates

$$\begin{split} h_1^0 &= \frac{1}{\sqrt{2}} \left(h_{dR}^0 + h_{uR}^0 \right) \qquad M_{h_1^0}^2 &= \frac{\lambda^2 v^2}{2} \\ h_2^0 &= \frac{1}{\sqrt{2}} \left(h_{dR}^0 - h_{uR}^0 \right) \qquad M_{h_2^0}^2 &= M_Z^2 + \frac{1}{2} \lambda^2 \left(2 v_S^2 - v^2 \right) \\ M_{s_R}^2 &= \frac{a_k v_S}{\sqrt{2}} + 2 \lambda^2 v_S^2 \end{split}$$

Two distinct phenomenologically viable scenarios occur.

- 1. $M_{h_1^0}^2 < M_{h_2^0}^2 \Rightarrow h = h_1^0$, $H = h_2^0 \Rightarrow \alpha = -\pi/4 = \beta \pi/2$. *H* does not have any direct tree level coupling to *WW* and *ZZ* that explains why the LHC does not see presently any other resonance but the lightest one at 125 GeV.
- 2. $M_{h_1^0}^2 > M_{h_2^0}^2 \Rightarrow h = h_2^0$, $H = h_1^0 \Rightarrow \alpha = \pi/4 = \beta$. LHC observed the second heaviest CP-even state because the couplings of the lightest one to fermions and to gauge bosons are strongly suppressed. Discovering such a light "sterile" Higgs boson is very difficult at the LHC.

Stable minimum conditions

We must ensure that all physical square masses are positive, which is equivalent to checking that our solution is a minimum of the potential. Moreover the constraint on $M_{H^{\pm}}^2$ implies that we are not breaking the $U(1)_{\rm em}$. So such a requirement will impose further constraints on the free parameters, that can be summarized as follows:

$$\operatorname{sign}(a_k) = -\operatorname{sign}(v_5), \quad |a_k| < 2\sqrt{2}\lambda^2 |v_5|$$

and one of the two following options

a) $v_{5}^{2} > \frac{1}{2}v^{2}$, b) $\frac{1}{2}\lambda^{2}(v^{2} - 2v_{5}^{2}) < M_{W}^{2}$.

From now on we shall assume that the lightest CP even scalar is the one coupled to the W's, thus $M_{h_1^0}^2 < M_{h_2^0}^2$. Moreover we want also $M_{h_1^0}^2 > M_Z^2$. This implies that the only available option is **a**).

Higgs production. 7 TeV



- 4 回 2 - 4 回 2 - 4 回 2 - 4

Э

Heavy chargino double degeneracy

$$M_{\tilde{c}} = \begin{pmatrix} \mathbf{0} & \mathbf{X}^{T} \\ \mathbf{X} & \mathbf{0} \end{pmatrix} \qquad \mathbf{X} = \begin{pmatrix} M_{2} & M_{W} \\ M_{W} & \mu \end{pmatrix} \qquad M_{H^{\pm}}^{2} \to \mu$$

$$M_{\chi_{L,H}^{2}}^{2} = \text{Eigenvalues}(\mathbf{X}^{T}\mathbf{X}) = \frac{1}{2} \begin{pmatrix} \mu^{2} + M_{2}^{2} + 2M_{W}^{2} \pm (\mu + M_{2})\sqrt{(M_{2} - \mu)^{2} + 4M_{W}^{2}} \end{pmatrix}$$

$$\downarrow$$
Solve $M_{\chi_{L}^{+}}^{2}$ in function of M_{2} : 2nd degree eq. $\to 2$ sols
$$\downarrow$$

$$M_{2} = \frac{M_{\chi_{L}^{+}}^{4} - M_{W}^{2} \mp \mu M_{\chi_{L}^{+}}^{2}}{\pm M_{\chi_{L}^{-}}^{2} - \mu}$$

$$\downarrow$$

$$M_{\chi_{H}^{2}}^{2} = \frac{\left(\mu(\mu \pm M_{\chi_{L}^{+}}^{2}) + M_{W}^{2}\right)^{2}}{(\mu \pm M_{\chi_{L}^{+}}^{2})^{2}}$$

Signal rates 8 TeV



Antonio Racioppi Fermiophobic Higgs boson and supersymmetry

イロン イヨン イヨン イヨン

Э

BR



Antonio Racioppi Fermiophobic Higgs boson and supersymmetry

・ロト ・回 ト ・ヨト ・ヨト

æ